
Early Treatment of Hyperdivergent Open-Bite Malocclusions

*Peter H. Buschang, MA, PhD, Wayne Sankey, DDS, MS, and
Jeryl D. English, DDS, MS*

This article establishes the morphologic attributes that characterize hyperdivergent open-bite (Hyp-OB) subjects, reviews associated etiologic factors, compares the various treatment modalities, and provides a rationale and a possible approach for early treatment. Treatment of Hyp-OB patients must address three-dimensional dentoalveolar and skeletal problems in both jaws. For early treatment to be successful, it must effectively deal with the etiology of the problem and the resulting mandibular skeletal dysmorphology. Of the various treatment approaches, including high-pull headgear, extractions, and bite blocks, the vertical chin cup holds the greatest potential for mandibular skeletal modification. Although early treatment can be theoretically justified based on psychosocial benefits and growth potential, more clinical and experimental research is required to optimize the treatment approach and define the long-term consequences. (Semin Orthod 2002;8:130-140.) Copyright 2002, Elsevier Science (USA). All rights reserved.

Treatment is ultimately dependent on an appropriate diagnosis, which in turn requires an unambiguous description of the problem. Before the advent of cephalometrics, orthodontists necessarily focused on dental relationships and defined open-bite malocclusion based on the vertical relationships of the maxillary and mandibular teeth. By the early 1960s, it became evident that distinctions needed to be made between open-bite malocclusions that included a skeletal component and those that did not.

Simple open-bite malocclusions that do not include skeletal components present less of a challenge for orthodontists; such open-bites of-

ten correct spontaneously.^{1,2} Because spontaneous correction occurs in up to 80% of mixed-dentition open-bite cases, it has been suggested that interceptive treatments are of little or no value.³ Successful treatment approaches are also available for persistent cases that do not self-correct.^{4,5}

In contrast, open-bite malocclusions that include skeletal components have proven to be extremely challenging for orthodontists. Most investigators have simply referred to them as skeletal open bites^{6,7}; Schudy⁸ characterized them as hyperdivergent, which reflects the skeletal phenotype.

Morphologic Characteristics

The typical hyperdivergent open-bite (Hyp-OB) patient presents with three-dimensional skeletal and dentoalveolar problems pertaining to both the maxilla and mandible. Variation in expression of traits among subjects should be expected. Within-subject differences clearly show that most of the dysmorphology occurs in the mandible.

Table 1 summarizes the problem list for the maxilla. The most consistent findings across

From the Department of Orthodontics, Baylor College of Dentistry, The Texas A&M University System Health Science Center, Dallas, TX; Private Practice, Flower Mound, TX; and Department of Orthodontics, University of Texas Health Science Center Houston, Houston, TX.

Address correspondence to Peter H. Buschang, Department of Orthodontics, Baylor College of Dentistry, The Texas A&M University System Health Science Center, 3302 Gaston Avenue, Dallas, TX 75246.

*Copyright 2002, Elsevier Science (USA). All rights reserved.
1073-8746/02/0803-0004\$35.00/0
doi:10.1053/sodo.2002.125432*

Table 1. Maxillary Morphologic Characteristics of Untreated Open Bites

Author	Heights		Length	SNA	PPA	Dentoalveolar Heights		Width
	Ant	Post				Incisor	Molar	
Subtelny and Sakuda ⁶	0	0	0	?	0	↑	↑	?
Bell ¹⁰⁵	?	?	?	?	?	↑	?	↓
Isaacson et al ¹⁰⁶	0	?	?	↓	?	↑	↑	↓
Nahoum et al ¹⁰⁷	↓	?	↓	?	↓	0	0	?
Schendel et al ¹⁰⁸	0	?	?	↓	?	↑	?	?
Lundström and Woodside ¹⁰⁹	↓	0	?	?	?	?	?	?
Fields et al ¹¹⁰	0	0	?	0	?	↑	↑	?
Nanda ^{95,96}	↓	0	?	?	↓	?	?	?
Haralabakis et al ¹¹¹	0	0	↓	?	?	↑	↑	0
Trouten et al ¹¹²	0	↑	?	?	↓	?	?	?
Lopez-Gavito et al ¹¹³	↓	0	?	↓	↓	↑	↑	?
Janson et al ¹¹⁴	?	0	?	?	?	↑	↑	?

Abbreviations: ↑, increase; ↓, decrease; 0, no difference; ?, not reported.

studies pertain to anterior and posterior dentoalveolar heights, which tend to be excessive. When evaluated, the palatal plane angles have been reported to be flatter because of decreased anterior upper facial heights; posterior upper facial heights do not appear to be affected. There is also a tendency for maxillas to be shorter and slightly recessive. When transverse dimensions have been evaluated, the maxilla is often reported to be narrow with an increased incidence of posterior cross-bites.

The most consistent mandibular characteristics are increased lower anterior facial heights, steeper mandibular planes, and larger gonial angles (Table 2). These traits combine to produce increased lower to upper and lower to total anterior facial height ratios. Most studies also

report decreased posterior facial heights caused by smaller ramus heights. It has also been shown that the glenoid fossa is positioned more superior relative to sella turcica. Mandibular dentoalveolar heights are also generally reported to be excessive in Hyp-OB patients.

Etiology of Hyp-OB Malocclusion

It is well established that abnormal muscles and habits are associated with Hyp-OB malocclusion.⁶ Finger, thumb, and tongue habits are perhaps the best known physical factors that produce open-bite malocclusions by lowering the mandible and preventing normal eruption.⁹ Individuals with prolonged nonnutritive-sucking habits have been repeatedly shown to have a

Table 2. Mandibular Morphologic Characteristics of Untreated Open Bites

Author	Heights		Corpus Lt	Gonial Angle	MPA	Retrogn	Dentalveolar Heights	
	Ant	Post					Incisor	Molar
Schudy ⁸	↑	↓	?	?	↑	↑	?	↑
Subtelny and Sakuda ⁶	↑	↓	0	↑	↑	↑	0	0
Bell ¹⁰⁵	↑	↓	?	↑	↑	↑	↑	?
Isaacson et al ¹⁰⁶	↑	↓	?	?	↑	↑	?	↑
Nahoum et al ¹⁰⁷	↑	↓	↓	↑	↑	?	0	↓
Schendel et al ¹⁰⁸	↑	↓	?	?	↑	↑	?	↑
Lundström and Woodside ¹⁰⁹	↑	?	?	↑	↑	↑	?	?
Cangialosi ¹¹⁵	↑	↓	?	↑	↑	?	?	?
Fields et al ¹¹⁰	↑	↓	0	↑	↑	0	↑	↑
Nanda ^{95,96}	↑	0	?	↑	?	?	?	?
Haralabakis et al ¹¹¹	?	↓	↓*	↑	?	?	↑	0
Trouten et al ¹¹²	?	?	?	↑	↑	?	?	?
Lopez-Gavito et al ¹¹³	↑	↑	?	?	↑	↑	0	0
Janson et al ¹¹⁴	?	?	?	?	?	?	↑	↑

Abbreviations: ↑, increase; ↓, decrease; 0, no difference; ?, not reported.

*Females only.

decrease overbite,¹⁰⁻¹⁴ increased overjet,^{10,11,13} decreased palatal width,^{11,15} and increased lip incompetence.¹²

There is considerable evidence showing smaller, less active, muscles and weaker bite forces among hyperdivergent subjects. Masseter and medial pterygoid muscle volumes and cross-sectional areas have been positively correlated with posterior face and ramal heights and negatively correlated with the mandibular plane and gonial angles.¹⁶⁻²¹ Kiliaridis and Kalebo²² showed that women with thin masseter muscles had proportionately longer faces. Bakke et al²³ found that muscle thickness at the most voluminous, superficial portion of the masseter was significantly correlated with bite force, anterior face height, and mandibular inclination. Moreover, the amplitudes of masseter and temporalis electromyographic (EMG) activity at rest, during swallowing, chewing, and maximal biting have also been negatively related with mandibular hyperdivergence in children 9 to 11 years of age.²⁴ Bakke and Michler²⁵ showed that EMG activity during maximal voluntary contraction was negatively correlated with anterior face height, mandibular inclination, vertical jaw relation, and gonial angle.

The smaller, less active muscles that characterize Hyp-OB subjects produce lower than normal bite forces.²⁶ In adults, stronger molar and incisor bite forces are highly correlated with smaller gonial angles, lower mandibular plane angles (MPA), and larger posterior face heights.^{27,28}

The relationship between weaker bite forces and hyperdivergence is not as strong for children as adults. Proffit and Fields²⁹ originally showed no significant differences in occlusal forces between long-face and normal children 6 to 11 years of age. By using more refined analytical techniques, Garcia-Morales et al³⁰ recently found that children (age 9.3 ± 3.6 years) with greater skeletal hyperdivergence had poorer mechanical advantage and lower maximum bite forces.

Although the association is well established, it remains controversial whether lower occlusal forces in long-face individuals produce hyperdivergence²⁶ or whether weaker occlusal forces are a biomechanical result of the long vertical facial proportions.³¹ This controversy is at least partially resolved by studies showing typical Hyp-OB

features among individuals with genetically determined neuromuscular diseases that weaken masticatory muscles. Patients with myotonic dystrophy have 2 to 3 times less EMG activity of the temporalis and masseter muscles during maximum clenching, lower maximum bite forces, anterior open bite malocclusions, and hyperdivergent growth patterns.^{32,33} Similarly, individuals with spinal muscular atrophy have open-bite malocclusions, increased vertical skeletal dimensions, steep mandibular plane angles, and hyperdivergent facial patterns.³⁴ They also have weaker, less efficient, masticatory muscles that fatigue more quickly than those of matched controls.³⁵

Numerous experimental animal studies and human clinical studies have also shown relationships between the Hyp-OB phenotype and chronic upper, middle, and lower airway obstructions. Rhesus monkeys forced to become mouth breathers showed greater than expected increases in anterior face height associated with mandibular lowering, decreases in arch width, and changes in tongue position and shape.³⁶ In fact, both light and heavy nasal obstruction produces downward and backward mandibular rotation, increased gonial angulation, and anterior open bite in primates³⁷ and rodents.³⁸

The relationship between airway obstruction and skeletal hyperdivergence is well established in humans.^{39,40} Compared with their nose-breathing counterparts, chronically allergic mouth-breathing children 6 to 12 years of age have narrower maxillas; greater incidence of posterior cross-bites; longer anterior facial heights; steeper palatal, occlusal, and mandibular planes; larger gonial angles; and more retrognathic mandibles.⁴¹ Mouth breathers with hypertrophied adenoids have narrow maxillas, lower tongue positions, proclined incisors, and increased lower anterior facial heights.⁴² Normalization of incisor inclination, maxillary arch width, depth of the bony nasopharynx, and mandibular plane inclination has been reported for children after adenoidectomies to correct severe nasopharyngeal obstruction.^{43,44} Children with enlarged tonsils also have more retrognathic mandibles, larger lower anterior facial heights, and larger MPAs.⁴⁵ The controversy concerning airway relates more to our ability to accurately diagnose airway problems for individuals rather

Table 3. Outcomes of Various Treatments Modalities for Hyperdivergent Open-Bite Malocclusion

<i>Effect Site</i>	<i>HPHG</i>	<i>HPGH + Splint</i>	<i>Extract</i>	<i>HPHG + Extract</i>	<i>Passive PBB</i>	<i>Active PBB</i>	<i>VCC</i>
Condylar growth/amt	–	–	0	0	0	0	+
Condylar growth/dir	0	0	0	0	–	0	+
Mandibular AP Position	0	0	0	0	+	+	+
Posterior Face Height	–	–	0	0	+	0	+
Anterior Face Height	0	0	0	0	+	+	0
Skeletal AP Relations	+	+	0	0	+	+	+
U6 Position	++	++	–	++	+	+	+
L6 Position	–	0	–	–	+	+	+
Overbite	0	0	+	++	+++	+	+
Overjet	+	+	+	++	+	+	+

Abbreviations: 0, no difference; –, not significant; +, significant; ++, very significant; +++, extremely significant.

than in the relationship between obstruction and skeletal hyperdivergence.

The common denominator among Hyp-OB subjects with habits, weak muscles, and airway obstructions appears to be a lowered mandibular posture. An individual's response to airway obstruction is immediate and predictable; the activity of the neck and masticatory muscles, head posture, and mandibular posture change quickly.^{46,47} Habits and weak elevator muscles might also produce changes in mandibular posture that, during growth, necessitate developmental adaptations leading to the Hyp-OB dysmorphology.

Treatment Modalities

Based on the forgoing, treatment of Hyp-OB patients must address three-dimensional problems pertaining to the dentoalveolar and skeletal structures of both jaws. Although hyperdivergent Class III open-bite cases are perhaps the most difficult to treat nonsurgically, they occur less frequently. In contrast, the Class II is the most common type of hyperdivergent patient. A severe hyperdivergent Class II patient requires reductions in dentoalveolar height throughout the maxilla and mandible (Table 3); reductions in gonial angulation; increased palatal plane angulation; maxillary expansion; and true mandibular autorotation to increase posterior mandibular height, redirect condylar growth, decrease anterior lower facial height, and reposition the chin forward (Fig 1).

High-Pull Headgear

High-pull headgear (HPHG) has traditionally been the appliance of choice for treating

Hyp-OB patients because it has been shown to effectively hold maxillary sutural growth and vertical dentoalveolar development.⁴⁸⁻⁵³ Although animal studies have reported absolute distal and superior displacement of metallic implants in the maxilla,^{54,55} human studies do not support increased autorotation.^{49,56} In fact, Baumrind and coworkers^{51,57} found that high-pull patients displayed relative increases in the MPA and reduced condylar growth.

Acrylic splints with HPHG create one large anchor unit that prevents unfavorable tipping of the upper molars.^{54,55,58-61} Caldwell and collaborators⁶² showed that this approach produced a superior and distal displacement of the maxilla, reductions of the Sella-Nasion-A (SNA) angle, clockwise rotation of the palatal plane, and relative intrusion of the upper molars. Importantly, they also reported increased lower molar eruption, decreased mandibular growth, and increased Sella-Nasion-B (SNB) angulation.

Extractions

Extraction therapy for hyperdivergent patients is predicated on the belief that molars moved mesially out of the occlusal wedge increase mandibular autorotation, decrease anterior facial height, and reduce open-bite malocclusions. However, Yamaguchi and Nanda⁶³ reported no differences in molar position or the A point-Nasion-B (ANB) angulation between extraction and nonextraction patients treated with high-pull face bows. Stagers⁶⁴ showed no significant differences between Class I nonextraction and extraction cases in changes of anterior facial height, upper to lower face height ratios, MPAs, posterior to anterior facial height ratios, or the distances of the molars to the palatal and man-




Figure 1. Problem lists for open-bite malocclusions. (AFH = anterior facial height; ant = anterior; post = posterior; md = mandible; mx = maxilla)

dibular planes. After extractions, Class II patients show decreases in the ANB angle, supraeruption of the lower molars, and increases in the MPA.⁶⁵ Pearson⁶⁶ evaluating cases with moderately steep MPAs treated with extractions and occipital headgears also reported significant supraeruption of the lower molars. Clearly, forward mandibular rotation and loss or even maintenance of vertical facial and dentoalveolar heights does not occur in the extraction patients.

Combined HPHG and extraction treatment produces results similar to extractions only. The primary differences pertain to the molars; the vertical movements of the upper molar are better controlled, but the lower molar shows even greater compensatory supraeruption.^{65,66} The positive effects of combined HPHG and extraction treatment are confined to the maxillary dentoalveolar regions. If the goal is to improve the orientation, position, and shape of the mandible, then other treatment approaches may be required.

Posterior Bite Blocks

Posterior bite blocks (PBB) have been shown to effectively modify vertical skeletal patterns in an-

imal models⁶⁷⁻⁷³ and humans.⁷⁴⁻⁸¹ However, PBBs hinge the mandible open beyond its resting position by varying amounts, which tends to increase the gonial angle.^{69,72,79} McNamara⁷⁹ concluded that the maxillary complex was most affected by PBB, although changes have been reported to occur throughout the craniofacial complex.⁶⁷

Animal studies evaluating repelling magnets embedded in bite-block appliances show superior-anterior maxillary displacement and molar intrusion. However, they also show greater potential for root resorption, deviated mandibular jaw posture that could produce skeletal asymmetries, and lateral open bites.^{70,73,75,76} Kalra and Burstone,⁷⁶ evaluating fixed magnetic bite blocks, reported increased mandibular length, decreased facial convexity, intrusion of the lower and upper molars, improved overjet and molar relationships, and small decreases in the mandibular plane angle and y axis. After treatment, posterior tooth eruption closed the posterior open bite, and the transverse jaw deviations self-corrected. More pronounced treatment effects have been reported for magnetic than spring-loaded bite blocks.⁸⁰

Iskan and associates⁷⁴ compared the effects of

a spring-loaded bite block (SLBB) worn for 6 months to a passive bite block and a vertical chin cup worn for 8 months. Both groups showed similar amounts of forward maxillary displacement, increases in mandibular length, posterior molar intrusion, mandibular autorotation, increased overbite, and reductions of anterior facial height. The SLBB group showed greater reductions in the ANB angle and lower molar intrusion, whereas the passive bite block and vertical chin cup group showed greater improvement in lower facial height and overbite. The SLBB also showed gonial angle increases, an undesirable effect.

Dellinger's⁷⁷ active vertical corrector, with repelling magnets embedded in bite blocks and acrylic shields to prevent lateral jaw deviations, is used in conjunction with a vertical chin cup. Intrusion of the posterior teeth, mandibular autorotation, and reductions in anterior height have been shown after 4 to 7 months of treatment. Dellinger⁷⁸ has also reported good long-term stability for five treated cases. Barbre and Sinclair,⁸¹ evaluating the effects of the active vertical corrector without the vertical chin cup, showed increased overbite, decreased lower anterior facial height, and reductions in mandibular plane angulation associated with upper and lower molar intrusion.

Vertical Chin Cup

Pearson⁸²⁻⁸⁴ has used the vertical chin cup in the mixed and permanent dentition to reduce the mandibular plane angle and limit increases in anterior facial height. A case treated with a vertical-pull chin cup in conjunction with a Kloehn cervical headgear showed significant dental and skeletal alterations; upper molar eruption and descent of the maxilla were inhibited while mandibular growth was redirected toward a more horizontal direction.⁸⁵ Treatment success was attributed to the increase in posterior facial height. Chin cups have also been used during active rapid palatal expansion (RPE) therapy to minimize the vertical displacement of the maxilla and control the opening of the mandibular plane angle.⁸⁶ Importantly, the vertical chin cup is the only appliance shown to effectively alter mandibular shape by increasing posterior heights, redirecting condylar growth, and decreasing gonial angulation.

Timing of Treatment

Early treatment can be best justified based on psychosocial benefits and growth potential for correction. Because facial appearance is the most important determinant of physical appearance⁸⁷ and the oral region contributes most to overall facial appearance,⁸⁸ abnormalities in the oral region negatively affect interpersonal relations,⁸⁹ how individuals are perceived by others,^{90,91} and self-perception.⁹² Early treatment might also obviate the costs and risks associated with relatively complex two-jaw surgical procedures typically performed to correct Hyp-OB malocclusions.

In terms of growth potential, early treatment of Hyp-OB is predicated on the knowledge that (1) it can be diagnosed early, (2) the phenotype does not self-correct, (3) potential for mandibular rotation and associated remodeling is greatest during childhood, (4) certain characteristics may require long periods of growth to fully correct, and (5) treatments are available that can correct complex three-dimensional configuration of problems.

The Problem Is Apparent Early and Does Not Self-Improve

Longitudinal studies agree that the Hyp-OB phenotype develops early and, on average, does not worsen with age. Longitudinal comparisons show a strong tendency to maintain long, average, and short facial type; most individuals (77%) present with the same facial type at 5 and 25.5 years of age.⁹³ Although there are growth differences between open- and deep-bite subjects in anterior face height,⁹⁴ the overall pattern of development is established early, even before the eruption of the first permanent molars, and maintained during growth. The mandibular plane and gonial angles of both open- and deep-bite groups decrease with age, which tend to decrease the absolute magnitude of skeletal imbalance for open-bite subjects.⁹⁵ Untreated vertical growers show less increase of Sella-Nasion-Pogonion (S-N-Pg), less decrease in the MPA, and less decrease in the gonial angle than horizontal growers.⁹⁶ If treatment mechanics of long-face individuals require orthopedic modification, it has been suggested that treatment should be initiated before the adolescent spurt.

Based on various measures of vertical development, Garcia-Morales and Buschang⁹⁷ recently showed that untreated subjects classified as hyperdivergent at 6 years of age maintain their dysmorphology relative to average or hypodivergent subjects through 15 years of age (Fig 2). Correlations between the individuals' 6- and 15 year-old phenotypes ranged between 0.5 to 0.75. Approximately 64% of the 6-year-old subjects classified as having a high MPA were also classified with high MPAs at 15 years; 28% had average MPAs and 8% had low MPAs (Fig 3A). Changes that occurred were related to the 15-year-old phenotype (correlations range 0.5-0.7) but not to the 6-year-old phenotype. In other words, knowing what a young child looks like helps predict his/her adult status but not his/her growth changes. Although most 6-year-old subjects with high MPAs improved, 24% became more hyperdivergent (Fig 3B). Together, these data suggest that the apparent stability displayed between group averages should not be expected for individual patients. These data emphasize that early interceptive treatment of Hyp-OB patients cannot be justified because we cannot currently predict whose malocclusions will worsen.

Childhood as a Period of Greater Potential

Overall growth potential and the capacity of certain characteristics to change are greater during childhood than adolescence. It is also possible that younger children would be more cooperative and more willing to undergoing complex long-lasting treatment regimens than adolescents.

Longitudinal analyses show that boys with high angles display significantly less true or total




Figure 2. Future growth changes for 6-year-old children with low (<1 SD), medium (\pm 1 SD) and high (>1 SD) mandibular plane angles (S-N/Go-Me).




Figure 3. (A) Frequencies of subjects with high, medium, and low MPAs (S-N/Go-Me) at 6 and 15 years of age. (B) Frequency of MPA (S-N/Go-Me) increasing and decreasing between 6 to 15 years for subjects classified with high, medium and low MPA (S-N/Go-Me) at 6 years.

forward rotation than low angle boys, with the difference being most pronounced during childhood.⁹⁸ Changes in the MPA angle also show significant group differences during childhood but not during adolescence. Spady et al⁹⁹ showed that true mandibular rotation was significantly greater during childhood than adolescence; rotation was particularly marked during the transition from primary to early mixed dentitions.

As shown, Hyp-OB patients often present with excessive maxillary and mandibular dentoalveolar heights. Relative intrusion of the teeth, especially mandibular teeth that show less eruption potential than their maxillary counterparts, may require considerable time for extreme cases. For example, the average mandibular incisor erupts 5.0 mm for boys and 3.7 mm for girls between 8 to 15 years of age; the first molar erupts 0.5 to 0.7 mm more than the incisor.¹⁰⁰ Relative intrusion over extended periods of growth could serve as an important and potent treatment modality.

Early Treatment Results

It is becoming increasingly clear that treatments for Hyp-OB patients should be directed toward the mandible. Naumann et al¹⁰¹ have shown that mandibular skeletal changes were twice as important as mandibular dental changes and 2.5 times as important as maxillary changes in affecting overbite changes. Vertical mandibular growth was shown to be more important than mandibular rotation in determining overbite changes. Their models suggest using appliances that limit anterior vertical mandibular growth and augment its forward rotation, rather than appliances that restrict maxillary growth.

Complex problems require complex treatment approaches. Sankey and coworkers¹⁰² recently reported outcomes for patients treated with lip seal exercises, a lower Crozat/lip bumper, a bonded palatal expander constructed to serve as a bite block, and a high-pull chin cup. A sample of children 8.2 ± 1.2 years with average MPAs of $40.1^\circ \pm 1.2^\circ$ performed lip seal exercises for 60 consecutive minutes each day. The Crozat/lip bumper appliance was cemented in place with 2 to 3 mm activation (after 8 weeks, it was reactivated 1 mm every 8 weeks). At the same time, the upper arch was expanded slowly (1/4 turn per week, 1 mm per month) for approximately 6 months. The expander infringed on the freeway space approximately 2 to 3 mm and was ramped to produce progressively thicker occlusal coverage on the palatal half of the appliance. A high-pull chin cup (force directed approximately 45° to occlusal plane) delivering 16 to 20 oz was worn at least 14 hours per day. The patients were compared with controls matched for age, gender, and MPA.

After 1.3 ± 0.3 years of treatment, condylar growth had been increased and changed toward a more anterosuperior direction. Maxillary expansion did not increase vertical dimensions. The mandible showed almost three times more forward rotation than expected, posterior facial height increased significantly more, the molars showed relative intrusion, the articular angle increased, the gonial angle decreased, and the chin moved forward almost twice as much as in the controls. Overjet decreased and overbite increased, especially for the patients with severe open-bite malocclusions. The aggregate of changes implies corrections in all three planes

of space that combine to produce a remarkable treatment result. Interestingly, a subsample of patients that performed clenching exercises sufficient to produce marks on the expander's acrylic did not wear high-pull chin cups but showed similar treatment outcomes.

The potential importance of clenching exercise in the treatment of Hyp-OP patients should not be underestimated. Tran and coworkers¹⁰³ recently completed a prospective clinical trial evaluating the effects of light clenching exercises as adjunctive treatment for HPHG and rapid palatal expansion in young children. Although HPHG without exercise produced den-toalveolar effects in the maxilla only, HPHG in combination with exercises also increased true mandibular autorotation and produced significant reductions in the ANB and gonial angles. Because there was no evidence of increased muscle strength, it was suggested that exercise might have altered the postural position of the mandible, which in turn influenced its rotational and remodeling patterns.

Conclusions

Although early treatment of hyperdivergent open-bite cases is theoretically appealing and practically possible, it remains poorly understood and must be approached with caution. More clinical and experimental research is required to establish the psychosocial benefits and to precisely define the long-term consequences of early treatment. The stability of early treatment will undoubtedly be related to the orthodontist's ability to correct the original cause of the problem. To that end, open-mouth posture associated with habits, weak muscles, or respiratory obstructions must be eliminated to avoid relapse. This implies additional studies necessary to develop sensitive and specific diagnostic indicators. Growth is clearly a critical period that holds great potential for orthopedic and orthodontic corrections as well as for relapse toward the original condition.

References

1. Hellman M: Open bite. *Int J Orthod* 17: 421-436, 1931
2. Korkhaus G: The frequency of orthodontic anomalies at various ages. *Int J Orthod* 14:120-129, 1928
3. Worms FW, Meskin LH, Isaacson RJ: Open-bite. *Am J Orthod* 59:589-595, 1971

4. Parker JH: The interception of the open bite in the early growth period. *Angle Orthod* 41:24-44, 1971
5. Johnson ED, Larson BE: Thumb-sucking: Classification and treatment. *J Dent Child* 60:392-398, 1993
6. Subtelny JD, Sakuda M: Open-bite: Diagnosis and treatment. *Am J Orthod* 50:337-358, 1964
7. Sassouni V: A classification of skeletal facial types. *Am J Orthod* 55:109-123, 1969
8. Schudy FF: Vertical growth versus anteroposterior growth as related to function and treatment. *Angle Orthod* 34:75-92, 1964
9. Graber TM: Thumb and finger-sucking. *Am J Orthod* 45:257-264, 1959
10. Larrson E: Dummy- and finger-sucking habits with special attention to their significance for facial growth and occlusion. 4. Effects on facial growth and occlusion. *Sven Tandlak Tidsskr* 65:605-634, 1972
11. Hawkins AC: A constructive approach to thumbsucking habit. *J Clin Orthod* 12:846-848, 1978
12. Bowden DB: A longitudinal study of the effects of digit- and dummy-sucking. *Am J Orthod* 52:887-901, 1966
13. Melsen B, Stensgaard K, Pedersen J: Sucking habits and their influence on swallowing pattern and prevalence of malocclusion. *Eur J Orthod* 1:271-280, 1979
14. Nanda RS, Khan I, Anand R: Effect of oral habits on the occlusion in preschool children. *J Dent Child* 39:449-452, 1972
15. Willmot DR: Thumb sucking habit and associated dental differences in monozygotic twins. *Br J Orthod* 11: 195-199, 1984
16. Weijts WA, Hillen B: Relationships between muscle cross-section and skull shape. *J Dent Res* 63:1154-1165, 1984
17. Gionhaku N, Lowe A: Relationship between jaw muscle volume and craniofacial form. *J Dent Res* 68:805-809, 1989
18. Hannam AG, Wood WW: Relationships between the size and spatial morphology of human masseter and medial pterygoid muscles, the craniofacial skeleton, and jaw biomechanics. *Amer J Phys Anthropol* 80:429-445, 1989
19. Van Spronsen PH, Weijts WA, Valk J, et al: Relationships between jaw muscle cross-sections and craniofacial morphology in normal adults, studied with magnetic resonance imaging. *Eur Orthod Soc* 13:351-361, 1991
20. Van Spronsen PH, Weijts WA, Valk J, et al: A Comparison of jaw muscle cross-sections of long-face and normal adults. *J Dent Res* 71:1279-1285, 1992
21. Benington PC, Gardener JE, Hunt NP: Masseter muscle volume measured using ultrasonography and its relationship with facial morphology. *Eur J Orthod* 21:659-670, 1999
22. Kiliaridis S, Kalebo P: Masseter muscle thickness measured by ultrasonography and its relation to facial morphology. *J Dent Res* 70:1262-1265, 1991
23. Bakke M, Tuxen A, Vilmann P, et al: Ultrasound image of human masseter muscle related to bite force, electromyography, facial morphology, and occlusal factors. *Scand J Dent Res* 100:164-171, 1992
24. Ingervall B, Thilander B: Relation between facial morphology and activity of the masticatory muscles. *J Oral Rehab* 1:131-147, 1974
25. Bakke M, Michler L: Temporalis and masseter muscle activity in patients with anterior open bite and cranio-mandibular disorders. *Scand J Dent Res* 99:219-228, 1991
26. Proffit W, Fields H, Nixon W: Occlusal forces in normal- and long-face adults. *J Dent Res* 62:566-571, 1983
27. Ringqvist M: Isometric bite force and its relation to dimensions of the facial skeleton. *Acta Odont Scand* 31:35-42, 1973
28. Kiliaridis S, Johansson A, Haraldson T, et al: Craniofacial morphology, occlusal traits, and bite force in persons with advanced occlusal tooth wear. *Am J Orthod Dentofacial Orthop* 107:286-292, 1995
29. Proffit W, Fields H: Occlusal forces in normal and long-face children. *J Dent Res* 62:571-574, 1983
30. Garcia-Morales P, Buschang PH, Throckmorton GS, et al: Maximum bite force, muscle efficiency and mechanical advantage in children with vertical growth pattern. *Eur J Orthod* 2002
31. Throckmorton G, Finn R, Bell WH: Biomechanics of differences. I. Lower face height. *Am J Orthod* 77:410-420, 1980
32. Harvold EP, Vargervik K, Chierici G: Primate experiments on oral sensation and dental malocclusions. *Am J Orthod* 63:494-508, 1973
32. Odman C, Kiliaridis S: Masticatory muscle activity in myotonic dystrophy patients. *J Oral Rehab* 23:5-10, 1996
33. Kiliaridis S, Mejerso C, Thilander B: Muscle function and craniofacial morphology: a clinical study in patients with myotonic dystrophy. *Eur J Orthod* 11:131-138, 1989
34. Houston KD, Buschang PH, Iannaccone ST, et al: Craniofacial morphological characteristics in patients with spinal muscular atrophy. *Pediatric Res* 36:265-269, 1994
35. Granger M, Buschang P, Throckmorton G, et al: Masticatory muscle function in patients with spinal muscular atrophy. *Am J Orthod* 115:697-702, 1999
37. Yamada, T, Tanne K, Miyamoto K, et al: Influences of nasal respiratory obstruction on craniofacial growth in young *Macaca fuscata* monkeys. *Am J Orthod Dentofacial Orthop* 111:38-43, 1997
38. Scarano E, Ottaviani SE, DeGirolamo S, et al: Relationship between chronic nasal obstruction and craniofacial growth: an experimental model. *Int J Pediatr Otorhinolaryngol* 2:125-131, 1998
39. Subtelny JD: The significance of adenoid tissue in orthodontia. *Ang Orthod* 25:59-69, 1954
40. McNamara JA: Influence of respiratory pattern on craniofacial growth. *Angle Orthod* 51:269-300, 1981
41. Bresolin D, Shapiro PA, Shapiro GG, et al: Mouth breathing in allergic children: its relationship to dento-facial development. *Am J Orthod* 83:334-340, 1983
42. Linder-Aronson S: Adenoids: Their effect on mode of breathing and nasal airflow and their relationship to characteristics of the facial skeleton and the dentition. *Acta Otolaryngologica Suppl* 265:1-132, 1970
43. Linder-Aronson S: Respiratory function in relation to facial morphology and the dentition. *Brit J Orthod* 6:59-71, 1979
44. Woodside DG, Linder-Aronson S, Lundstrom A, et al:

- Mandibular and maxillary growth after changed mode of breathing. *Am J Orthod Dentofacial Orthop* 100:1-18, 1991
45. Behlfelt K, Linder-Aronson S, McWilliam J, et al: Cranio-facial morphology in children with and without enlarged tonsils. *Eur J Orthod* 12:233-243, 1990
 46. Hellsing E, Forsberg CM, Linder-Aronson S, et al: Changes in postural EMG activity in the neck and masticatory muscles following obstruction of the nasal airway. *Eur J Orthod* 8:247-253, 1986
 47. Tourne LP, Schweiger J: Immediate postural responses to total nasal obstruction. *Am J Orthod Dentofacial Orthop* 110:606-11, 1996
 48. Armstrong MM: Controlling the magnitude, direction, and duration of extraoral force. *Am J Orthod* 59:217-243, 1971
 49. Watson WG: A computerized appraisal of the high-pull facebow. *Am J Orthod* 62:561-573, 1972
 50. Badell M: An evaluation of extraoral combined high-pull traction and cervical traction to the maxilla. *Am J Orthod* 69:431-445, 1976
 51. Baumrind S, Korn, EL, Molthen, BS, et al: Changes in facial dimensions associated with the use of forces to retract the maxilla. *Am J Orthod* 80:17-30, 1981
 52. Baumrind S, Korn EL, Isaacson RJ, et al: Quantitative analysis of the orthodontic and orthopedic effects of maxillary traction. *Am J Orthod* 84:384-398, 1983
 53. Firouz M, Zernik J, Nanda R: Dental and orthopedic effects of high-pull headgear in treatment of Class II Division I malocclusion. *Am J Orthod* 102:197-205, 1992
 54. Elder JR, Tuenge RH: Cephalometric and histologic changes produced by high-pull traction to the maxilla in *Macaca mulatta*. *Am J Orthod* 66:599-617, 1974
 55. Meldrum RJ: Alterations in the upper facial growth of *Macaca mulatta* resulting from high-pull headgear. *Am J Orthod* 67:393-411, 1975
 56. Worms FW, Isaacson RJ, Speidel TM: A concept and classification of centers of rotation and extraoral force systems. *Angle Orthod* 43:384-401, 1973
 57. Baumrind S, Molthen R, West EE, et al: Mandibular plane changes during maxillary retraction. *Am J Orthod* 74:32-40, 1978
 58. Thurow RC: Craniomaxillary orthopedic correction with en masse dental control. *Am J Orthod* 68:601-624, 1975
 59. Joffe L, Jacobsen A: The maxillary orthopedic splint. *Am J Orthod* 75:54-69, 1979
 60. Fotis V, Melsen B, Williams S, et al: Vertical control as an important ingredient in the treatment of severe sagittal discrepancies. *Am J Orthod* 86:224-232, 1984
 61. Orton HS, Slattery DA, Orton S: The treatment of severe 'gummy' class II division I malocclusion using a maxillary intrusion splint. *Eur J Orthod* 14:216-223, 1992
 62. Caldwell SF, Hymas TA, Timm TA: Maxillary traction splint: A cephalometric evaluation. *Am J Orthod* 85:376-384, 1984
 63. Cangialosi TJ: Skeletal features of anterior open bite. *Am J Orthod* 85:28-36, 1984
 63. Yamaguchi K, Nanda RS: The effects of extraction and nonextraction treatment on the mandibular position. *Am J Orthod* 100:443-452, 1991
 64. Staggers JA: Vertical changes following first premolar extractions. *Am J Orthod Dentofacial Orthop* 105:19-24, 1994
 65. Dougherty HL: The effect of mechanical forces upon the mandibular buccal segments during orthodontic treatment. *Am J Orthod* 54:29-49, 1968
 66. Pearson LE: Vertical control through use of mandibular posterior intrusive forces. *Angle Orthod* 43:194-200, 1973
 67. Sergl HG, Farmand M: Experiments with unilateral bite planes in rabbits. *Angle Orthod* 45:108-114, 1975
 68. Altuna G: The effect of excess occlusal force on the eruption of the buccal segments and maxillary and mandibular growth direction in the *Macaca* monkey. Master's thesis, University of Toronto, Toronto, 1979
 69. Altuna G, Woodside DG: Response of the midface to treatment with increased vertical occlusal forces. Treatment and post-treatment effects in monkeys. *Angle Orthod* 55:251-263, 1985
 70. Woods MG, Nanda RS: Intrusion of posterior teeth with magnets—an experiment in growing baboons. *Angle Orthod* 58:136-150, 1988
 71. Woodside DG, Linder-Aronson S: Progressive increase in lower anterior facial height and the use of posterior occlusal bite-block in its management, in Graber LW (ed): *Orthodontics: State of the Art, Essence of the Science*, St Louis, MO, Mosby, 1986, pp 209-218
 72. Rowe, TK, Carlson, DS: The effect of bite-opening appliances on the mandibular rotational growth and remodeling in the rhesus monkey (*Macaca mulatta*). *Am J Orthod Dentofacial Orthop* 98:544-549, 1990
 73. Melsen B, McNamara JA, Hoenie DC: The effect of bite-blocks with and without repelling magnets studied histomorphologically in the rhesus monkey (*Macaca mulatta*). *Am J Orthod Dentofacial Orthop* 108:500-509, 1995
 74. Iscan HN, Akkaya S, Koralp E: The effects of the spring-loaded posterior bite block on the maxillo-facial morphology. *Eur J Orthod* 14:54-60, 1992
 75. Kiliardis S, Egermark I, Thilander B: Anterior open bite treatment with magnets. *Eur J Orthod* 13:447-457, 1990
 76. Kalra V, Burstone CJ: Effects of a fixed magnetic appliance on the dentofacial complex. *Am J Orthod Dentofacial Orthop* 95:467-478, 1989
 77. Dellinger EL: A clinical assessment of the active vertical corrector—A nonsurgical alternative for skeletal open-bite. *Am J Orthod Dentofacial Orthop* 89:428-436, 1986
 78. Dellinger EL: Active vertical corrector treatment—Long-term follow-up of anterior open bite treated by the intrusion of posterior teeth. *Am J Orthod Dentofacial Orthop* 110:145-154, 1996
 79. Lundström A, Woodside DG: Longitudinal changes in facial type in cases with vertical and horizontal mandibular growth directions. *Eur J Orthod* 5:259-268, 1983
 79. McNamara JA: An experimental study of increased vertical dimension in the growing face. *Am J Orthod* 71:382-395, 1977
 80. Kuster R, Ingervall B: The effect of treatment of skeletal open bite with two types of bite-blocks. *Eur J Orthod* 14:489-499, 1992

81. Barbre RE, Sinclair PM: A cephalometric evaluation of anterior open bite correction with the magnetic active vertical corrector. *Angle Orthod* 61:93-109, 1991
82. Pearson LE: Vertical control in treatment of patients having backward-rotational growth tendencies. *Angle Orthod* 48:132-140, 1978
83. Pearson LE: Vertical control in fully-banded orthodontic treatment. *Angle Orthod* 56:205-224, 1986
84. Pearson LE: Treatment of a severe openbite excessive vertical pattern with an eclectic non-surgical approach. *Angle Orthod* 61:71-76, 1991
85. Haas AJ: A biological approach to diagnosis, mechanics and treatment of vertical dysplasia. *Angle Orthod* 50:279-300, 1980
86. Majourau A, Nanda R: Biomechanical basis of vertical dimension control during rapid palatal expansion therapy. *Am J Orthod Dentofacial Orthop* 106:322-328, 1996
87. Alley TR: Social and applied aspects of face perception: An introduction, in Alley TR (ed): *Social and Applied Aspects of Perceiving Faces*. Hillsdale, NJ, Lawrence Erlbaum Assoc, 1988, pp 1-8
88. Berschied E, Walster E, Bohrnstedt G: The happy American body: A survey report. *Psychol Today* 7:119-131, 1973
89. Macgregor FC: Social and psychological implications of dentofacial disfigurement. *Angle Orthod* 40:231-233, 1970
90. Dion KK, Berscheid E, Walster E: What is beautiful is good. *J Pers Soc Psychol* 24:285-293, 1972
91. Shaw WC: The influence of children's dentofacial appearance on their social attractiveness as judged by peers and lay adults. *Am J Orthod* 79:399-415, 1981
92. Kiyak HA, McNeill RW, West RA, et al: Personality characteristics as predictors and sequelae of surgical and conventional orthodontics. *Am J Orthod Dentof Orthop* 89:383-392, 1986
93. Bishara SE, Jakobsen JR: Longitudinal changes in three normal facial types. *Am J Orthod* 88:466-502, 1985
94. Nanda SK: Patterns of vertical growth in the face. *Am J Orthod Dentofacial Orthop* 93:103-116, 1988
95. Nanda SK: Growth patterns in subjects with long and short faces. *Am J Orthod Dentofacial Orthop* 98:247-258, 1990
96. Lundström A, Woodside DG: Longitudinal changes in facial type cases with vertical and horizontal mandibular growth directions. *Eur J Orthod* 5:259-268, 1983
97. Garcia-Morales P, Buschang PH: Longitudinal stability of divergent growth patterns. *J Dent Res* 81:388, 2002
98. Karlson AT: Craniofacial growth differences between low and high MP-SN angle males: a longitudinal study. *Angle Orthod* 65:341-350, 1995
99. Spady M, Buschang PH, Demirjian A: Mandibular rotation and angular remodeling during childhood and adolescence. *Am J Human Biol* 4:683-689, 1992
100. Watanabe E, Demirjian A, Buschang PH: Longitudinal post-eruptive mandibular tooth movements of males and females. *Eur J Orthod* 21:459-468, 1999
101. Naumann SA, Behrents RG, Buschang PH: Vertical components of overbite change: A mathematical model. *Am J Orthod Dentofacial Orthop* 117:486-495, 2000
102. Sankey WL, Buschang PH, English J, et al: Early treatment of vertical skeletal dysplasia: The hyperdivergent phenotype. *Am J Orthod Dentofacial Orthop* 118:317-327, 2000
103. Tran MT, English JD, Throckmorton GS, et al: The adjunctive treatment effects of light masticatory muscle training on hyperdivergent open-bite patients. A pilot study. 2002 (in press)
104. Bell WH, Creekmore TD, Alexander RG: Surgical correction of the long face syndrome. *Am J Orthod* 71:40-67, 1977
105. Isaacson JR, Isaacson RJ, Speidel TM, et al: Extreme variation in vertical facial growth and associated variation in skeletal and dental relations. *Angle Orthod* 41:219-229, 1971
106. Nahoum HI, Horowitz SL, Benedicto EA: Varieties of anterior open-bite. *Am J Orthod* 61:486-492, 1972
107. Schendel SA, Eisenfeld J, Bell WH, et al: The long face syndrome: Vertical maxillary excess. *Am J Orthod* 70:398-408, 1976
108. Lundström A, Woodside DG: A comparison of various facial and occlusal characteristics in mature individuals with vertical and horizontal growth direction expressed at the chin. *Eur J Orthod* 3:227-235, 1981
109. Fields H, Proffit W, Nixon W, et al: Facial pattern differences in long-faced children and adults. *Am J Orthod* 85:217-223, 1984
110. Haralabakis NB, Yiagtzis SC, Toutountzakis NM: Cephalometric characteristics of open bite in adults: A three dimensional cephalometric evaluation. *Int J Adult Orthod Orthognath Surg* 9:223-231, 1994
111. Trouten JC, Enlow DH, Rabine M, et al: Morphologic factors in open bite and deep bite. *Angle Orthod* 53:192-211, 1983
112. Lopez-Gavito G, Wallen TR, Little R, et al: Anterior open-bite malocclusion: A longitudinal 10-year postretention evaluation of orthodontically treated patients. *Am J Orthod* 87:175-186, 1985
113. Janson GR, Metaxas A, Woodside DG: Variation in maxillary and mandibular molar and incisor vertical dimension in 12-year-old subjects with excess, normal, and short lower anterior facial height. *Am J Orthod* 106:409-418, 1994
114. Congialosi TJ: Skeletal morphologic features of anterior open-bite. *Am J Orthod* 85:28-36, 1984